

Infection control for technologists and radiologists

Herein we illustrated an example, but we encourage you consulting the experts in your hospital or local authorities.

Standard Precaution



Patient: Wear a surgical mask

Medical personnel:

Use Personal Protective Equipment (PPE), i.e surgical mask, apron, gloves, and face shield/goggle.

When there is potential for aerosol transmission:

Use an N95 mask.

Don't forget to perform fit-check ahead of time



#1. Hand hygiene



- Use enough amount of hand sanitizer
- Cover all surfaces until hands feel dry
- ▶ This should take around 20-30 seconds



#2 Wear a long-sleeved gown





#3 Put on a surgical mask



Use an N95 mask when necessary



#4 Wear gloves





#5 Cover the couch with a sheet of disposable drape





#6 Call in the patient (Let the patient wear a surgical mask)



The patient does not necessarily have to change clothes.



#1. Take off the PPE before entering the operation room










#2. Hand hygiene





#3 Take the gown off and discard it into the medical waste container






Treat the front side of PPE as being contaminated.



#4. Hand hygiene





#5 Take off the mask(s)







#6. Hand hygiene



And now you are ready to come back to the operation room.

#1. Hand hygiene




Scan finished!



#2 Wear a long-sleeved gown



#3 Put on a surgical mask




#4 Hand hygiene and then wear gloves





#1 Take off the PPE and discard them in a medical waste container







#2. After the hand hygiene, wear apron and gloves.





#3. Discard the drape





#4. Clean the surface where patient might have touched





#5. Clean the doorknobs both inside and outside of the room.





Do not forget cleaning the panels on CT scanner.



#6. Discard the PPE into the medical waste container.





#7. Air change if necessary.



For COVID-19, air change is considered unnecessary unless there is potential for creating aerosol in the room.

That's it!



Produced by the volunteers of JCR




















> Supervision: Naohisa Fujita MD PHD, Kyoto Prefectural University of Medicine

